

COMC 2012 Official Solutions

by Adrian Tang (adrian.l.tang@gmail.com)

A1 Determine the positive integer n such that $8^4 = 4^n$.

Solution: The answer is $n = 6$.

Solution 1: Note that $8^4 = (2^3)^4 = 2^{12} = 4^6$. Therefore, $n = 6$. $\square$

Solution 2: We write 8^4 and 4^n as an exponent with base 2.

$$\begin{aligned} 8^4 &= 4^n \\ (2^3)^4 &= (2^2)^n \\ 2^{12} &= 2^{2n} \end{aligned}$$

Therefore, $2n = 12$. Hence, $n = 6$. $\square$

Solution 3: Note that $8^4 = (8^2)^2 = 64^2 = 4096$. Hence, $4^n = 4096$. We check each positive integer n starting from 1.

n	4^n
1	4
2	16
3	64
4	256
5	1024
6	4096

All positive integers $n > 6$ yield a value of 4^n larger than 4096. Therefore, $n = 6$. $\square$

- A2 Let x be the *average* of the following six numbers: $\{12, 412, 812, 1212, 1612, 2012\}$. Determine the value of x .

Solution: The answer is $x = 1012$.

Solution 1: The sum of the first and sixth terms is 2024. The sum of the second and fifth terms is 2024 and the sum of the third and fourth terms is 2024. Hence, the sum of the six terms is 2024×3 . Hence, the average of the six terms is

$$\frac{2024 \times 3}{6} = \frac{2024}{2} = 1012. \quad \square$$


Solution 2: The average of the six numbers is

$$\begin{aligned} & \frac{12 + 412 + 812 + 1212 + 1612 + 2012}{6} \\ &= \frac{0 + 400 + 800 + 1200 + 1600 + 2000}{6} + \frac{12 + 12 + 12 + 12 + 12 + 12}{6} \\ &= \frac{100(4 + 8 + 12 + 16 + 20)}{6} + 12 = \frac{100 \times 60}{6} + 12 = 1000 + 12 = 1012. \quad \square \end{aligned}$$

Solution 3: Note that the sequence is arithmetic.¹ Therefore, the average of the six numbers is the average of the middle two numbers, which is the halfway point between 812, 1212. Hence, the answer is 1012. $\square$


¹A sequence is said to be arithmetic if successive terms in the sequence have a common difference.

- A3 Let $ABCDEF$ be a hexagon all of whose sides are equal in length and all of whose angles are equal. The area of hexagon $ABCDEF$ is exactly r times the area of triangle ACD . Determine the value of r .



Solution 1: The answer is $r = 3$.

Divide the hexagon into six regions as shown, with the centre point denoted by P .



This is possible since the hexagon is regular. By symmetry, note that $PA = PB = PC = PD = PE = PF$ and the six interior angles about P are equal. Then since the sum of the six interior angles about P sum to 360° ,

$$\angle APB = \angle BPC = \angle CPD = \angle DPE = \angle EPF = \angle FPA = 60^\circ.$$


Therefore, the six triangles $\triangle PAB, \triangle PBC, \triangle PCD, \triangle PDE, \triangle PEF, \triangle PFA$ are all equilateral and have the same area. Let K be the area of any one of these triangles. Therefore, the hexagon has area $6K$.

Note that the area of $\triangle ACD$ is equal to the area of $\triangle PCD$ plus the area of $\triangle PAC$. Since $\triangle PAB, \triangle PBC$ are both equilateral, $PA = AB$ and $PC = CB$. Therefore, triangles $\triangle BAC$ and $\triangle PAC$ are congruent and hence have the same area. Note that the area of $\triangle PAC$ plus the area of $\triangle BAC$ is the sum of the areas of the equilateral triangles $\triangle PAB$ and $\triangle PBC$, which is $2K$. Therefore, $\triangle PAC$ has area K . We already noted that the area of $\triangle ACD$ is

equal to the area of $\triangle PCD$ plus that of $\triangle PAC$. This quantity is equal to $K + K = 2K$. Hence, the area of $ABCDEF$ is $6K/2K = 3$ times the area of $\triangle ACD$. The answer is 3. $\square$

Solution 2: Divide the hexagon into six regions and define K as in Solution 1. Note that $\triangle APC$ and $\triangle DPC$ have a common height, namely the height from C to AD . Since $PA = PD$, $\triangle APC$ and $\triangle DPC$ have the same area, namely K . Therefore, the area of $\triangle ACD$ is the sum of the areas of $\triangle APC$ and that of $\triangle DPC$, which is $K + K = 2K$. Hence, the ratio of the area of $ABCDEF$ to the area of $\triangle ACD$ is $6K/2K = 3$. $\square$

Solution 3: The sum of the angles of a hexagon is $180^\circ \times (6 - 2) = 720^\circ$. Therefore, $\angle ABC = 120^\circ$. Since $BA = BC$, $\angle BAC = \angle BCA$. Then since the sum of the angles of $\triangle ABC$ is 180° and $\angle ABC = 120^\circ$, $\angle BAC = \angle BCA = 30^\circ$. Since $\angle BCA = 30^\circ$ and $\angle BCD = 120^\circ$, $\angle ACD = 90^\circ$.



Suppose that each side of the hexagon has length 1. We now determine the length AC to determine the area of $\triangle ACD$. By the cosine law,

$$AC^2 = BA^2 + BC^2 - 2 \cdot BA \cdot BC \cdot \cos \angle ABC = 1^2 + 1^2 - 2 \cdot 1 \cdot 1 \cdot \cos 120^\circ = 2 - 2 \cdot (-1/2) = 3.$$

Therefore, $AC = \sqrt{3}$. Hence, the area of $\triangle ACD$ is $1/2 \cdot CD \cdot CA = 1/2 \cdot 1 \cdot \sqrt{3} = \sqrt{3}/2$.

We now find the area of the hexagon. As in Solution 1, the hexagon consists of 6 equilateral triangles each with side 1. The area of each equilateral triangle is $\sqrt{3}/4$. Therefore, the area of the hexagon is $6 \cdot \sqrt{3}/4 = 3\sqrt{3}/2$. Therefore, the ratio of the area of the hexagon to the area of $\triangle ACD$ is

$$\frac{3\sqrt{3}/2}{\sqrt{3}/2} = 3.$$

Therefore, the answer is 3. $\square$

Solution 4: As in Solution 3, suppose each side of the hexagon has length 1. Then the area of $\triangle ACD$ is $\sqrt{3}/2$. Note that the area of $\triangle ABC$ is


$$\frac{1}{2} \cdot BA \cdot BC \cdot \sin 120^\circ = \frac{1}{2} \cdot 1 \cdot 1 \cdot \frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{4}.$$

Therefore, $\triangle ACD$ is twice the area of $\triangle ABC$ and then $\triangle ACD$ is $2/3$ the area of the quadrilateral $ABCD$. But the line AD splits the hexagon $ABCDEF$ in half. Therefore, $\triangle ACD$ is $1/3$ the area of the entire hexagon. Therefore, the ratio of the area of the hexagon to the area of $\triangle ACD$ is 3. $\square$

Solution 5: We will follow Solution 4, but provide a different way to show that the area of $\triangle ACD$ is twice the area of $\triangle ABC$. Note that these two triangles have a common height with base AD and BC , respectively. Since AD is twice the length of BC , $\triangle ACD$ is twice the area of $\triangle ABC$. Then as in Solution 4, we can conclude that the ratio of the area of the hexagon to the area of $\triangle ACD$ is 3. $\square$

Solution 6: Join the segment DF . The hexagon is cut into four triangles. By symmetry, $\triangle ACD$ and $\triangle AFB$ are congruent, as are $\triangle ABC$ and $\triangle DEF$. Note that $AD \parallel BC$ and $FC \parallel BF$. Let AD, FC meet at P . Then $\triangle APC$ and $\triangle ABC$ are congruent (parallelogram cut by diagonal). $\triangle APC$ has half the height of $\triangle ABC$ on base AC (by symmetry), so $[ACD] = 2[ABC]$, where $[\dots]$ denotes the area of a figure. Similarly, $[AFD] = 2[ABC]$. Thus the hexagon's area is $[ACD] + [ADF] + [ABC] + [DEF] = 6[ABC] = 3[ACD]$. Therefore, the answer is 3. $\square$

- A4 Twelve different lines are drawn on the coordinate plane so that each line is parallel to exactly two other lines. Furthermore, no three lines intersect at a point. Determine the total number of intersection points among the twelve lines.



Solution: The answer is 54.

Solution 1: Since no point lies on three or more lines, the number of intersection points is equal to the number of pairs of lines that intersect. The total number of pairs of lines is $12 \times 11/2 = 6 \times 11 = 66$. Each line is parallel to two other lines. Hence, each line is part of two pairs of lines that do not intersect. Since there are twelve lines, there are $12 \times 2/2 = 12$ pairs of lines that do not intersect. Therefore, there are $66 - 12 = 54$ pairs of lines that intersect. Hence, the answer is 54. $\square$

Solution 2: Since each line is parallel to exactly two other lines, each line is not parallel to nine other lines. Hence, each line intersects nine other lines.


Since no point lies on three or more lines, each point of intersection lies on exactly two lines. Combining these two observations yields that the total number of intersection points is

$$\frac{12 \times 9}{2} = 54. \quad \square$$

Solution 3: Recall that two lines are parallel if and only if they have the same slope. Consider the set of slopes of the twelve lines; since each line is parallel to exactly two other lines, each slope is the slope of three lines among the twelve lines. Hence, there are four different slopes represented among the twelve lines.

Since each slope contains three lines, each pair of slopes contains $3 \times 3 = 9$ points of intersection. There are four different slopes. Hence, the number of pairs of different slopes is $4 \times 3/2 = 6$. Since no three lines intersect at a common point, the number of points of intersection is $9 \times 6 = 54$. $\square$

- B1 Alice and Bob run in the clockwise direction around a circular track, each running at a constant speed. Alice can complete a lap in t seconds, and Bob can complete a lap in 60 seconds. They start at diametrically-opposite points.



When they meet for the first time, Alice has completed exactly 30 laps. Determine all possible values of t .

Solution: The answer is $t = 59$ or $t = 61$.

Since Alice ran exactly 30 laps, Bob meets Alice at where Alice started. Since Bob started diametrically across from Alice, Bob ran $n + \frac{1}{2}$ laps for some positive integer n . Since Alice and Bob meet only the first time they meet, the number of laps that Alice ran and the number of laps Bob ran cannot differ by more than 1. Therefore, Bob ran either 29.5 laps or 30.5 laps.

Note that Alice and Bob ran for the same amount of time and the number of seconds each person ran is the number of laps he/she ran times the number of seconds it takes he/she to complete a lap.

If Bob ran 29.5 laps, then $30t = 29.5 \times 60$. Hence, $t = 29.5 \times 2 = 59$.

If Bob ran 30.5 laps, then similarly, $30t = 30.5 \times 60$. Hence, $t = 30.5 \times 2 = 61$.

Therefore, $t = 59$ or $t = 61$. $\square$

- B2 For each positive integer n , define $\varphi(n)$ to be the number of positive divisors of n . For example, $\varphi(10) = 4$, since 10 has 4 positive divisors, namely $\{1, 2, 5, 10\}$.

Suppose n is a positive integer such that $\varphi(2n) = 6$. Determine the minimum possible value of $\varphi(6n)$.

Solution: The answer is 8.

Solution 1: Recall that if a positive integer m has prime factorization $p_1^{e_1} p_2^{e_2} \dots p_t^{e_t}$, where $p_1, \dots, p_t$ are distinct primes, then the number of positive divisors of m is $\varphi(m) = (e_1 + 1)(e_2 + 1) \dots (e_t + 1)$ (*). Note that each term in this product is at least 2.

Since $2n$ is an even positive integer with 6 positive divisors, $2n = 2^5, 2^2 \cdot p$ or $2 \cdot p^2$, where p is some odd prime number. Therefore, $n = 2^4 = 16, 2p$ or p^2 . Therefore, $6n = 6 \times 16 = 96, 12p$ or $6p^2$.

Note that $\varphi(96) = \varphi(2^5 \times 3^1) = 6 \times 2 = 12$.

If $p = 3$, then $\varphi(12p) = \varphi(36) = \varphi(2^2 \times 3^2) = 3 \times 3 = 9$ and $\varphi(6p^2) = \varphi(54) = \varphi(2^1 \times 3^3) = 2 \times 4 = 8$.

It remains to show the case when $p > 3$. So far the minimum value obtained for $\varphi(6n) = 8$. If $p > 3$, then $6n$ contains at least 3 different prime divisors. Then by (*), the number of positive divisors of $6n$ is at least $2 \times 2 \times 2 = 8$. Therefore, $\varphi(6n) \geq 8$ for all positive integers n . As we have shown, $n = 9$ yields $\varphi(6n) = 8$. Therefore, the answer is 8. $\square$

Solution 2: Note that four positive divisors of $2n$ are $1, 2, n$ and $2n$. Note also that $n = 2$ does not satisfy $\varphi(2n) = 6$. Therefore, $n \geq 2$ and consequently, $1, 2, n$ and $2n$ are all distinct.

Since $2n$ has 6 positive divisors, there are two other positive divisors a, b of $2n$, with $a, b > 2$. Then the set of positive divisors of $2n$ is $\{1, 2, a, b, n, 2n\}$.

Now consider the positive divisors of $6n$. Note that the set of positive divisors of $6n$ contains those of $2n$. Further note that $3n$ and $6n$ are positive divisors of $6n$, which are not positive divisors of $2n$. Hence, the set of positive divisors of $6n$ contains $\{1, 2, a, b, n, 2n, 3n, 6n\}$. Therefore, $\varphi(6n) \geq 8$.

We will show that this minimum can be obtained. Since $\{1, 2, a, b, n, 2n, 3n, 6n\}$ are the positive divisors of $6n$ and appear in increasing order, $a \cdot 2n = 6n$ and $bn = 6n$. Multiplying both


equations and dividing both sides by $2n^2$ yield $ab = 18$. But since $\{1, 2, a, b, n, 2n\}$ are the positive divisors of $2n$, $ab = 2n$. Therefore, $2n = 18$, from which we can conclude that $n = 9$ is a candidate which yields $\varphi(2n) = 6$ and $\varphi(6n) = 8$.

This can be easily verified, since the positive divisors of 18 are $\{1, 2, 3, 6, 9, 18\}$. Since the positive divisors of 54 are $\{1, 2, 3, 6, 9, 18, 27, 54\}$, $\varphi(54) = 8$. $\square$

- B3 Given the following 4 by 4 square grid of points, determine the number of ways we can label ten different points $A, B, C, D, E, F, G, H, I, J$ such that the lengths of the nine segments

$$AB, BC, CD, DE, EF, FG, GH, HI, IJ$$

are in strictly increasing order.



Solution: The answer is 24.

First, we count the number of possible lengths of the segments. By the Pythagorean Theorem, the different lengths are $\sqrt{0^2 + 1^2} = 1, \sqrt{0^2 + 2^2} = 2, \sqrt{0^2 + 3^2} = 3, \sqrt{1^2 + 1^2} = \sqrt{2}, \sqrt{1^2 + 2^2} = \sqrt{5}, \sqrt{1^2 + 3^2} = \sqrt{10}, \sqrt{2^2 + 2^2} = \sqrt{8}, \sqrt{2^2 + 3^2} = \sqrt{13}, \sqrt{3^2 + 3^2} = \sqrt{18}$. These nine lengths are all different. Therefore, all nine lengths are represented among $AB, BC, CD, DE, EF, FG, GH, HI, IJ$. Furthermore, these nine lengths in increasing order are:


$$\begin{aligned} \sqrt{0^2 + 1^2} &< \sqrt{1^2 + 1^2} < \sqrt{0^2 + 2^2} < \sqrt{1^2 + 2^2} < \sqrt{2^2 + 2^2} \\ &< \sqrt{0^2 + 3^2} < \sqrt{1^2 + 3^2} < \sqrt{2^2 + 3^2} < \sqrt{3^2 + 3^2}. \end{aligned}$$

Hence, the longest length must be a segment that goes from one corner to the diagonally-opposite corner.


We will construct the ten points in the order $J, I, H, G, F, E, D, C, B, A$.

For simplicity, we place the points on the coordinate plane, with the bottom left corner at $(0, 0)$ and the top right corner at $(3, 3)$.


Note that J must be a corner of the grid, and there are four such corners. Furthermore, I must be the diagonally opposite corner from J . Without loss of generality, suppose $J = (0, 0)$. Then $I = (3, 3)$.




The point H has the property that $HI = \sqrt{3^2 + 2^2}$, i.e. H is a point which is distance three horizontally from I and distance two vertically from I , or vice versa. By symmetry along the diagonal JI , there two choices for H , namely $(0, 1)$ or $(1, 0)$. Without loss of generality, suppose $H = (1, 0)$.




The segment GH has length $\sqrt{3^2 + 1^2}$. Hence, G is either $(0, 3)$ or $(2, 3)$. But if $G = (0, 3)$ then F is a point such that $FG = 3 = \sqrt{0^2 + 3^2}$. Then $F = (0, 0)$ or $(3, 3)$, which are already occupied by J, I , respectively. Therefore, G cannot be $(0, 3)$, and thus must be $(2, 3)$. Consequently, $F = (2, 0)$.



EF has length $\sqrt{8} = \sqrt{2^2 + 2^2}$. Hence, $E = (0, 2)$. DE has length $\sqrt{5} = \sqrt{2^2 + 1^2}$. Hence, $D = (2, 1)$.




Then $C = (0, 1)$ and $B = (1, 2)$.



From B , there are three remaining points A such that $AB = 1$, namely $(1, 1), (1, 3), (2, 2)$.


By our construction, the points J, H and A were the only points where there was more than one choice. Every other point was determined from our construction. There were 4 choices for J , 2 choices for H and 3 choices for A . Hence, the number of ways to select 10 points that satisfy the condition given in the problem is $4 \times 3 \times 2 = 24$. The answer is 24. $\square$

- B4 In the following diagram, two lines that meet at a point A are tangent to a circle at points B and C . The line parallel to AC passing through B meets the circle again at D . Join the segments CD and AD . Suppose $AB = 49$ and $CD = 28$. Determine the length of AD .



Solution 1: The answer is $AD = 63$.

Join the segment BC . Since the two lines are both tangent to the circle, $AB = AC$. Therefore, $\angle ABC = \angle ACB$.



Furthermore, since BD is parallel to AC , $\angle ACB = \angle DBC$. Since AC is tangent to the circle at C , by the tangent-chord theorem, $\angle BDC = \angle ACB$. Hence, we have the following sequence of equal angles:


$$\angle ABC = \angle ACB = \angle CBD = \angle CDB.$$

Furthermore, $AB = AC$ and $CB = CD$. Therefore, $\triangle ABC$ is similar to $\triangle CBD$. Hence,

$$\frac{AB}{BC} = \frac{CB}{BD}.$$

Since $AB = 49$ and $BC = CD = 28$, $BD = BC^2/AB = 28^2/49 = 4^2 = 16$.

Let M be the foot of the perpendicular from D on AC and N the foot of the perpendicular on BD from C .



Since $CB = CD$, N is the midpoint of BD . Since BD is parallel to CM , $NDMC$ is a rectangle. Therefore, $CM = ND = \frac{1}{2} \cdot BD = 8$. We now determine the length of DM .


$$DM = NC = \sqrt{DC^2 - DN^2} = \sqrt{28^2 - 8^2} = \sqrt{784 - 64} = \sqrt{720}.$$

Therefore,

$$AD = \sqrt{AM^2 + MD^2} = \sqrt{(AC + CM)^2 + MD^2} = \sqrt{(49 + 8)^2 + 720} = \sqrt{3969} = 63.$$

Hence, the answer is 63. $\square$

Solution 2: As in Solution 1, join segment BC and conclude that $BC = 28$ and $BD = 16$. Also as in Solution 1, $\angle ABC = \angle ACB = \angle CBD = \angle CDB$. Let θ be this angle.



Let N be the foot of the perpendicular on BD from C . As in Solution 1, N is the midpoint of BD . Therefore, $BN = 8$. We can now determine $\cos \theta$ from $\triangle CBN$, which is

$$\cos \theta = \frac{BN}{BC} = \frac{8}{28} = \frac{2}{7}.$$

(Alternatively, we can use $\triangle ABC$ to determine $\cos \theta$.) Note that $\angle ABD = 2\theta$. We then apply the cosine law on $\triangle ABD$ to determine AD . By the cosine law, we have

$$AD^2 = BA^2 + BD^2 - BA \cdot BD \cdot \cos(2\theta).$$

We first determine $\cos(2\theta)$, which by the double-angle formula is

$$\cos(2\theta) = 2\cos^2\theta - 1 = 2 \cdot (2/7)^2 - 1 = -41/49.$$

Therefore,

$$AD^2 = 49^2 + 16^2 - 2(49)(16)(-41/49) = 2401 + 256 + 2 \cdot 16 \cdot 41 = 3969.$$

Hence, $AD = \sqrt{3969} = 63$. $\square$

Solution 3: Let θ be defined as in Solution 2. Then as shown in Solution 2, $\cos \theta = 2/7$. Then note that

$$\angle BCD = 180 - \angle CBD - \angle CDB = 180 - 2\theta.$$

Therefore, $\angle ACD = \angle ACB + \angle BCD = \theta + (180 - 2\theta) = 180 - \theta$. We now apply the cosine law on $\triangle ACD$.

$$\begin{aligned} AD^2 &= CA^2 + CD^2 - 2 \cdot CA \cdot CD \cdot \cos \angle ACD = 49^2 + 28^2 - 2 \cdot 49 \cdot 28 \cdot \cos(180 - \theta) \\ &= 2401 + 784 + 2 \cdot 49 \cdot 28 \cdot \cos \theta = 3185 + 2 \cdot 49 \cdot 28 \cdot \frac{2}{7} = 3185 + 4 \cdot 7 \cdot 28 = 3969. \end{aligned}$$

Therefore, $AD = \sqrt{3969} = 63$. $\square$

Part C

C1 Let $f(x) = x^2$ and $g(x) = 3x - 8$.

- (a) (2 marks) Determine the values of $f(2)$ and $g(f(2))$.
- (b) (4 marks) Determine all values of x such that $f(g(x)) = g(f(x))$.
- (c) (4 marks) Let $h(x) = 3x - r$. Determine all values of r such that $f(h(2)) = h(f(2))$.

Solution:

- (a) The answers are $f(2) = 4$ and $g(f(2)) = 4$.

Substituting $x = 2$ into $f(x)$ yields $f(2) = 2^2 = 4$.

Substituting $x = 2$ into $g(f(x))$ and noting that $f(2) = 4$ yields $g(f(2)) = g(4) = 3 \cdot 4 - 8 = 4$. $\square$

- (b) The answers are $x = 2$ and $x = 6$.

Note that

$$f(g(x)) = f(3x - 8) = (3x - 8)^2 = 9x^2 - 48x + 64$$

and

$$g(f(x)) = g(x^2) = 3x^2 - 8.$$

Therefore, we are solving

$$9x^2 - 48x + 64 = 3x^2 - 8.$$

Rearranging this into a quadratic equation yields

$$6x^2 - 48x + 72 = 0 \Rightarrow 6(x^2 - 8x + 12) = 0.$$

This factors into $6(x - 6)(x - 2) = 0$. Hence, $x = 2$ or $x = 6$. We now verify these are indeed solutions.

If $x = 2$, then $f(g(2)) = f(3(2) - 8) = f(-2) = (-2)^2 = 4$ and $g(f(2)) = 4$ by part(a). Hence, $f(g(2)) = g(f(2))$. Therefore, $x = 2$ is a solution.

If $x = 6$, then $f(g(6)) = f(3 \cdot 6 - 8) = f(10) = 10^2 = 100$ and $g(f(6)) = g(6^2) = g(36) = 3 \cdot 36 - 8 = 108 - 8 = 100$. Hence, $f(g(6)) = g(f(6))$. Therefore, $x = 6$ is also a solution. $\square$

(c) The answers are $r = 3$ and $r = 8$.

We first calculate $f(h(2))$ and $h(f(2))$ in terms of r .

$$f(h(2)) = f(3 \cdot 2 - r) = f(6 - r) = (6 - r)^2$$

and

$$h(f(2)) = h(2^2) = h(4) = 3 \cdot 4 - r = 12 - r.$$

Therefore, $(6 - r)^2 = 12 - r \Rightarrow r^2 - 12r + 36 = 12 - r$. Re-arranging this yields

$$r^2 - 11r + 24 = 0,$$

which factors as

$$(r - 8)(r - 3) = 0.$$

Hence, $r = 3$ or $r = 8$. We will now verify that both of these are indeed solutions.

If $r = 3$, then $h(x) = 3x - 3$. Then $f(h(2)) = f(3 \cdot 2 - 3) = f(3) = 9$ and $h(f(2)) = h(2^2) = h(4) = 3 \cdot 4 - 3 = 9$. Therefore, $f(h(2)) = h(f(2))$. Consequently, $r = 3$ is a solution. From the result of part (b), we also verified that $r = 8$ is a solution. $\square$

C2 We fill a 3×3 grid with 0s and 1s. We score one point for each row, column, and diagonal whose sum is *odd*.

1	1	0
1	0	1
0	1	1

1	1	1
1	0	1
0	1	1

For example, the grid on the left has 0 points and the grid on the right has 3 points.

(a) (2 marks) Fill in the following grid so that the grid has exactly 1 point. No additional work is required. Many answers are possible. You only need to provide one.

Solution: Any of the following is a solution:

0	0	0
1	1	0
1	1	0

1	1	0
1	1	0
0	0	0

0	1	1
0	1	1
0	0	0

0	0	0
0	1	1
0	1	1

0	1	1
1	1	0
1	0	1

1	0	1
1	1	0
0	1	1

1	0	1
0	1	1
1	1	0

1	1	0
0	1	1
1	0	1

(b) (4 marks) Determine all grids with exactly 8 points.

Solution: Note that there are three rows, three columns and two diagonals. Hence, every row, column and diagonal has an odd sum.

We will consider two cases; the first case is when the middle number is 0 and second case is when the middle number is 1.

Case 1: If the middle number is 0, then let A, B, C, D be the values provided in the following squares.

A	B	C
	0	D

Then since each row, column and diagonal has an odd sum, each term diametrically opposite from A, B, C, D has a different value from A, B, C, D , respectively. Denote $\bar{0} = 1$ and $\bar{1} = 0$. Then we have the following values in the grid:

A	B	C
$\bar{D}$	0	D
$\bar{C}$	$\bar{B}$	$\bar{A}$

Note that $X + \bar{X} = 1$ for any value X . Then note the sum of $A, B, C, \bar{A}, \bar{B}, \bar{C}$ is $1 + 1 + 1 = 3$. Hence, one of $A + B + C$ and $\bar{A} + \bar{B} + \bar{C}$ is even. Therefore, either the top row or bottom row sum to an even number. Hence, there are no grids with 8 points in this case.

Case 2: If the middle number is 1, then again, let A, B, C, D be the values provided in the following squares.

A	B	C
	1	D

Then since each row, column and diagonal has an odd sum, each term diagonally opposite from A, B, C, D has the same value as A, B, C, D , respectively. Then we have the following values in the grid:

A	B	C
D	1	D
C	B	A

Since $A + B + C$ and $A + D + C$ are both odd, $B = D$.

A	B	C
B	1	B
C	B	A

Hence, the only remaining restriction is that $A + B + C$ is odd. Since $A, B, C = 0$ or 1 , $A + B + C = 1$ or 3 . The only triples (A, B, C) that give this result are $(A, B, C) = (1, 0, 0), (0, 1, 0), (0, 0, 1)$ or $(1, 1, 1)$. The following are the grids corresponding to these results, which completes the problem. $\square$

1	0	0
0	1	0
0	0	1

0	1	0
1	1	1
0	1	0

0	0	1
0	1	0
1	0	0

1	1	1
1	1	1
1	1	1

- (c) (4 marks) Let E be the number of grids with an even number of points, and O be the number of grids with an odd number of points. Prove that $E = O$.

Solution 1: Consider the set of all grids. Pair the grids so that each grid G is paired with the grid G^* formed by switching the top-left number of G . (By switching, we mean if the top left number of G is 0, we switch it to a 1. If the top left number of G is 1, we switch it to a 0.) The following is an example of the action provided by G^* .

$$G = \begin{array}{|c|c|c|} \hline 0 & 0 & 1 \\ \hline 1 & 1 & 1 \\ \hline 1 & 0 & 1 \\ \hline \end{array} \quad G^* = \begin{array}{|c|c|c|} \hline 1 & 0 & 1 \\ \hline 1 & 1 & 1 \\ \hline 1 & 0 & 1 \\ \hline \end{array}$$

Note that the sum of the elements in the top row, the left most column and the diagonal going from the top-left to the bottom-right switches parity, i.e. switches either from odd to even, or even to odd and the sum of the elements of the other rows / columns / diagonals remain unchanged. Hence, the total number of rows/columns/diagonals which have odd sum in G and G^* differ by an odd number. Hence, exactly one of G, G^* has an even number of points and the other has an odd number of points. Since each grid lies in exactly one pair, there is the same number of grids with an even number of points as grids with an odd number of points, i.e. $E = O$. $\square$

Comment: The solution also applies if we switch any one of the four corners of the grid.

Solution 2: Note that the grid consisting of all zeros has an even number of points, namely zero. Note that for any grid, switching the centre square keeps the parity of the number of points the same. Switching any of the four side squares keeps the parity of the number of points the same. As in Solution 1, switching the centre changes the parity of the number of points the same.

Therefore, if a grid has 0, 2 or 4 of its corners as 1, then the number of points of the grid is even. If a grid has 1 or 3 of its corner as 1, then number of points in the grid is odd.


We will count the number of grids of based on the number of corner squares containing 1.

There are five non-corner squares. Therefore, there are 2^5 grids with zero corners containing 1.

There are $\binom{4}{1} = 4$ ways to choose one corner to be 1. Therefore, there are 4×2^5 grids with one corner containing 1. Similarly, there are $\binom{4}{2} \times 2^5 = 6 \times 2^5$ grids with two corners containing 1, $\binom{4}{3} \times 2^5 = 4 \times 2^5$ grids with three corners containing 1 and 2^5 grids with four corners containing 1.


Hence, there are $2^5(1 + 6 + 4) = 8 \times 2^5$ grids with an even number of points and $2^5(4 + 4) = 8 \times 2^5$ grids with an odd number of points. Therefore, $E = O$, as desired. $\square$

- C3 Let $ABCD$ be a parallelogram. We draw in the diagonal AC . A circle is drawn inside $\triangle ABC$ tangent to all three sides and touches side AC at a point P .



- (a) (2 marks) Prove that $DA + AP = DC + CP$.

Solution: Let the circle inside $\triangle ABC$ touch AB, BC at X, Y , respectively.



Then by equal tangents, we have

$$DA + AP = DA + AX = DA + AB - BX$$

and

$$DC + CP = DC + CY = DC + CB - BY.$$

By equal tangents, we have $BX = BY$. Since opposite sides of a parallelogram have equal lengths, $AB = DC$ and $DA = CB$. Therefore, $DA + AB - BX = DC + CB - BY$. Consequently, $DA + AP = DC + CP$, as desired. $\square$


- (b) (4 marks) Draw in the line DP . A circle of radius r_1 is drawn inside $\triangle DAP$ tangent to all three sides. A circle of radius r_2 is drawn inside $\triangle DCP$ tangent to all three sides. Prove that

$$\frac{r_1}{r_2} = \frac{AP}{PC}.$$


Solution 1: Consider the triangles $\triangle APD$ and $\triangle CPD$ and note that the heights of these triangles to side AP, PC are the same. Therefore,

$$\frac{AP}{PC} = \frac{[APD]}{[CPD]},$$

where $[\dots]$ denotes area.



Given any triangle QRS with a circle on the inside touching all three sides, let O be the centre of the circle and r the radius of the circle. Then the distance from O to each of the sides QR, RS, SQ is the same, and is the radius of the circle. Join OQ, OR, OS .



Then

$$\begin{aligned}
 [QRS] &= [OQR] + [ORS] + [OSQ] = \frac{r \cdot QR}{2} + \frac{r \cdot RS}{2} + \frac{r \cdot SQ}{2} \\
 &= \frac{r}{2} \cdot (QR + RS + SQ) = \frac{r}{2} \cdot (\text{Perimeter of } \triangle QRS).
 \end{aligned}$$

Then

$$[APD] = \frac{r_1}{2} \cdot (\text{Perimeter of } \triangle APD)$$

and

$$[CPD] = \frac{r_2}{2} \cdot (\text{Perimeter of } \triangle CPD)$$


Then

$$\frac{AP}{PC} = \frac{[APD]}{[CPD]} = \frac{r_1}{r_2} \cdot \frac{\text{Perimeter of } \triangle APD}{\text{Perimeter of } \triangle CPD}.$$

Hence, to prove that $AP/PC = r_1/r_2$, it suffices to show that $\triangle APD, \triangle CPD$ have the same perimeter.

By part (a), we have $DA + AP = DC + CP$. The perimeter of $\triangle APD$ is $DA + AP + PD = DC + CP + PD$, which is the perimeter of $\triangle CPD$. This solves the problem. $\square$

Solution 2: Let X, Y, Z be the centres of the circles inside $\triangle ABC$, $\triangle APD$ and $\triangle CPD$, respectively, M the point where the circle inside $\triangle ADP$ touch AC and N the point where the circle inside $\triangle CDP$ touch AC . Note that XP, YM and ZN are each perpendicular to AC .



Note also that AY bisects $\angle DAC$, CZ bisects $\angle DCA$, AX bisects $\angle BAC$ and CX bisects $\angle BCA$. Since AD is parallel to BC , $\angle DAC = \angle BCA$. Therefore, $\angle CA Y = \angle ACX$, which implies that $\angle MAY = \angle PCX$. Since $\triangle AYM$ and $\triangle CXP$ are both right-angled triangles, $\triangle AYM \sim \triangle CXP$. Similarly, $\triangle CZN \sim \triangle AXP$. Therefore,

$$\frac{AM}{MY} = \frac{CP}{PX}, \quad \text{and} \quad \frac{CN}{NZ} = \frac{AP}{PX}.$$


Note that $MY = r_1$ and $NZ = r_2$. This yields

$$\frac{AM}{r_1} = \frac{CP}{PX}, \quad \text{and} \quad \frac{CN}{r_2} = \frac{AP}{PX}.$$

Dividing the second equation by the first equation yields

$$\frac{AP}{PC} = \frac{AM}{CN} \cdot \frac{r_1}{r_2}.$$

Therefore, to solve the problem, it suffices to show that $AM = CN$.




Let the circle inside $\triangle DAP$ touch AD, DP at R, S , respectively. Then note that $AR = AM, DR = DS$ and $PM = PS$. Therefore, $DA + AP = DR + RA + AM + MP = DS + AM + AM + SP = 2AM + DP$. Similarly, $DC + CP = 2CN + DP$. By part (a), $DA + AP = DC + CP$. Therefore, $2AM + DP = 2CN + DP$, from which we can conclude that $AM = CN$. This solves the problem. $\square$

- (c) (4 marks) Suppose $DA + DC = 3AC$ and $DA = DP$. Let r_1, r_2 be the two radii defined in (b). Determine the ratio r_1/r_2 .

Solution: The answer is $r_1/r_2 = 4/3$.

Solution 1: By part (b). $r_1/r_2 = AP/PC$. Let $x = AP$ and $y = PC$. The answer is the ratio x/y .

By part (a), $DA + AP = DC + CP$. Let $s = DA + AP = DC + CP$. Then $DA = s - x$ and $DC = s - y$. Since $DA + DC = 3AC$, $(s - x) + (s - y) = 3(x + y)$. Hence, $2s = 4(x + y)$. Therefore, $s = 2(x + y)$. Therefore, $DA = x + 2y$ and $DC = 2x + y$. Since $DP = DA$, $DP = x + 2y$.



Drop the perpendicular from D to AC and let the perpendicular intersect AC at M . Since $DA = DP$, M is the midpoint of AP . Therefore, $MP = x/2$. By the Pythagorean Theorem, we have $MD^2 + MC^2 = DC^2$ and $MD^2 + MP^2 = DP^2$. Therefore, $DC^2 - MC^2 = DP^2 - MP^2$. Therefore,

$$(2x + y)^2 - (x/2 + y)^2 = (x + 2y)^2 - (x/2)^2.$$

Simplifying this yields

$$4x^2 + 4xy + y^2 - \frac{x^2}{4} - xy - y^2 = x^2 + 4xy + 4y^2 - \frac{x^2}{4}.$$

Hence, $3x^2 - xy - 4y^2 = 0$. Factoring this yields $(3x - 4y)(x + y) = 0$. Since x, y are lengths, $x + y \neq 0$. Therefore, $3x - 4y = 0$. Therefore, $x/y = 4/3$. $\square$

Solution 2: We define x, y as in Solution 1. Then we have $DA = x + 2y$ and $DC = 2x + y$ and $DP = x + 2y$. Consider triangles $\triangle ADP$ and $\triangle CDP$. Then

$$\cos \angle APD = \frac{PA^2 + PD^2 - AD^2}{2 \cdot PA \cdot PD} = \frac{x^2 + (x + 2y)^2 - (x + 2y)^2}{2 \cdot x \cdot (x + 2y)} = \frac{x^2}{2x(x + 2y)} = \frac{x}{2(x + 2y)}$$

and

$$\cos \angle CPD = \frac{PC^2 + PD^2 - CD^2}{2 \cdot PC \cdot PD} = \frac{y^2 + (x + 2y)^2 - (2x + y)^2}{2 \cdot y \cdot (x + 2y)} = \frac{-3x^2 + 4y^2}{2y(x + 2y)}.$$

Since $\angle APD$ and $\angle CPD$ sum to 180° , their cosine values are negatives of each other. Hence,

$$\frac{-x}{2(x + 2y)} = \frac{-3x^2 + 4y^2}{2y(x + 2y)} \Rightarrow -x = \frac{-3x^2 + 4y^2}{y}.$$

This simplifies to $3x^2 - xy - 4y^2 = 0$. Factoring this yields $(3x - 4y)(x + y) = 0$. As in Solution 1, we get $x/y = 4/3$. $\square$

Solution 3: We define x, y as in Solution 1. Then we have $DA = x + 2y$ and $DC = 2x + y$ and $DP = x + 2y$. We now determine $\cos \angle DAP$ using cosine law in both $\triangle DAP$ and $\triangle DAC$.

$$\begin{aligned} \cos \angle DAP &= \frac{AD^2 + AP^2 - DP^2}{2 \cdot AD \cdot AP} \\ &= \frac{(x + 2y)^2 + x^2 - (x + 2y)^2}{2 \cdot (x + 2y) \cdot x} = \frac{x^2}{2x(x + 2y)} = \frac{x}{2(x + 2y)} \end{aligned}$$

and

$$\begin{aligned} \cos \angle DAC &= \frac{AD^2 + AC^2 - DC^2}{2 \cdot AD \cdot AC} \\ &= \frac{(x + 2y)^2 + (x + y)^2 - (2x + y)^2}{2 \cdot (x + 2y)(x + y)} = \frac{-2x^2 + 2xy + 4y^2}{2(x + 2y)(x + y)} = \frac{-(x - 2y)(x + y)}{(x + 2y)(x + y)} = \frac{-x + 2y}{x + 2y}. \end{aligned}$$

Therefore,

$$\frac{x}{2(x + 2y)} = \frac{-x + 2y}{x + 2y}.$$

Hence, $x = 2(-x + 2y)$. This simplifies to $x/y = 4/3$. $\square$

C4 For any positive integer n , an n -tuple of positive integers $(x_1, x_2, \dots, x_n)$ is said to be *super-squared* if it satisfies both of the following properties:

- (1) $x_1 > x_2 > x_3 > \dots > x_n$.
- (2) The sum $x_1^2 + x_2^2 + \dots + x_k^2$ is a perfect square for each $1 \leq k \leq n$.

For example, $(12, 9, 8)$ is super-squared, since $12 > 9 > 8$, and each of 12^2 , $12^2 + 9^2$, and $12^2 + 9^2 + 8^2$ are perfect squares.

- (a) (2 marks) Determine all values of t such that $(32, t, 9)$ is super-squared.

Solution: The only answer is $t = 24$.

Note that $32^2 + t^2 = 1024 + t^2$ and $32^2 + t^2 + 9^2 = 1105 + t^2$ are perfect squares. Then there exist positive integers a, b such that

$$\begin{aligned} 1024 + t^2 &= a^2 \\ 1105 + t^2 &= b^2. \end{aligned}$$

Subtracting the first equation from the second equation gives

$$b^2 - a^2 = 81 \Rightarrow (b - a)(b + a) = 81.$$

The only ways 81 can be written as the product of two distinct positive integers is $81 = 1 \times 81$ and $81 = 3 \times 27$.

If $(b - a, b + a) = (1, 81)$, then $b - a = 1$ and $b + a = 81$. Summing these two equations yield $2b = 82$. Therefore, $b = 41$. Hence, $a = 40$. Therefore, $t^2 = a^2 - 32^2 = 40^2 - 32^2 = 8^2(5^2 - 4^2) = 8^2 \cdot 3^2$. Hence, $t = 24$.

We now verify that $(32, 24, 9)$ is indeed super-squared. Clearly, the tuple is strictly decreasing, i.e. satisfies condition (1). Finally, $32^2 + 24^2 = 8^2(4^2 + 3^2) = 8^2 \cdot 5^2 = 40^2$ and $32^2 + 24^2 + 9^2 = 40^2 + 9^2 = 1681 = 41^2$. Therefore, the tuple also satisfies condition (2).

If $(b - a, b + a) = (3, 27)$, then $b - a = 3$ and $b + a = 27$. Summing these two equations gives $2b = 30$. Therefore, $b = 15$. Hence, $a = 12$. Therefore, $t^2 = a^2 - 32^2 = 12^2 - 32^2 < 0$. Hence, there are no solutions for t in this case.

Therefore, $t = 24$ is the only solution.

- (b) (2 marks) Determine a super-squared 4-tuple (x_1, x_2, x_3, x_4) with $x_1 < 200$.

Solution: Note that if $(x_1, \dots, x_n)$ is super-squared, then $(ax_1, \dots, ax_n)$ is also super-squared for any positive integer a . We will show that this tuple satisfies both (1) and (2) to show that it is indeed super-squared. Clearly, since $x_1 > x_2 > \dots > x_n$, $ax_1 > ax_2 > \dots > ax_n$. Since $x_1^2 + x_2^2 + \dots + x_k^2$ is a perfect square, $x_1^2 + x_2^2 + \dots + x_k^2 = m^2$ for some positive integer m . Therefore, $(ax_1)^2 + \dots + (ax_k)^2 = (am)^2$. Hence, $(ax_1, \dots, ax_n)$ is super-squared.

From the example in the problem statement, $(12, 9, 8)$ is super-squared. Therefore, $12(12, 9, 8) = (144, 108, 96)$ is also super-squared. Note that $12^2 + 9^2 + 8^2 = 17^2$. Hence, $144^2 + 108^2 + 96^2 = 204^2 = 12^2 \cdot 17^2$.

Note that $13^2 \cdot 17^2 = (12^2 + 5^2) \cdot 17^2 = 12^2 \cdot 17^2 + 5^2 \cdot 17^2 = 12^2 \cdot 17^2 + 85^2$. Therefore, $221^2 = 13^2 \times 17^2 = 144^2 + 108^2 + 96^2 + 85^2$. And so we conclude that $(144, 108, 96, 85)$ is super-squared.

Comment: The list of all super-squared 4-tuples (x_1, x_2, x_3, x_4) with $x_1 < 200$ is

$$(132, 99, 88, 84), (144, 108, 75, 28), (144, 108, 96, 85), (156, 117, 104, 60), (180, 96, 85, 60), \\ (180, 135, 120, 32), \quad \text{and} \quad (192, 144, 100, 69).$$

(c) (6 marks) Determine whether there exists a super-squared 2012-tuple.

Solution: There does indeed exist a super-squared 2012-tuple.

We will show that there exists a super-squared n -tuple for any positive integer $n \geq 3$. We will prove this by induction on n . In the problem statement and in part (b), we showed that this statement holds for $n = 3, 4$.

Suppose there exists a super-squared k -tuple $(x_1, x_2, \dots, x_k)$ for some positive integer $k \geq 3$. We will show from this k -tuple that there exists a super-squared $(k+1)$ -tuple.

Let a, b, c be a tuple of positive integers such that $a^2 + b^2 = c^2$. We will provide the additional conditions on (a, b, c) shortly.

Let r be the positive integer such that $x_1^2 + x_2^2 + \dots + x_k^2 = r^2$. As in part (b), we note that if $(x_1, \dots, x_k)$ is super-squared, then $(ax_1, \dots, ax_k)$ is also super-squared and $(ax_1)^2 + \dots + (ax_k)^2 = (ar)^2$. Then we claim that $(ax_1, \dots, ax_k, br)$ satisfies property (2) of super-squared. Clearly, $(ax_1)^2 + \dots + (ax_t)^2$ is a perfect square, since $(ax_1, \dots, ax_k)$ is super-squared, for all $1 \leq t \leq k$. To prove the claim, it remains to show that $(ax_1)^2 + \dots + (ax_k)^2 + (br)^2$ is a perfect square. This is clear since this quantity is equal

to $(ar)^2 + (br)^2 = r^2(a^2 + b^2) = (cr)^2$. This proves the claim.

To make the tuple $(ax_1, \dots, ax_k, br)$ super-squared, we require that $ax_k > br$, or equivalently, $a/b > r/x_k$. Note that r, x_k are determined from the tuple $(x_1, \dots, x_k)$. Hence, it suffices to show that there exists a Pythagorean triple (a, b, c) , with $a^2 + b^2 = c^2$ such that $a/b > r/x_k$. In general, we need to show that a/b can be arbitrarily large.

Note that $(a, b, c) = (m^2 - 1, 2m, m^2 + 1)$ is a Pythagorean triple for any positive integer m . This is clear since $(m^2 - 1)^2 + (2m)^2 = m^4 - 2m^2 + 1 + 4m^2 = m^4 + 2m^2 + 1 = (m^2 + 1)^2$. In such a case,

$$\frac{a}{b} = \frac{m^2 - 1}{2m} = \frac{m}{2} - \frac{1}{2m} > \frac{m}{2} - 1,$$

which can be made arbitrarily large. This completes the induction proof. $\square$